






Wybrane algorytmy

-  Borkowski, K. M. (1989). “Accurate algorithms to transform geocentric to geodetic coordinates”. *Bulletin géodésique* 63.1, s. 50–56.
-  Borkowski, K. M. (1987). “Transformation of geocentric to geodetic coordinates without approximations”. *Astrophysics and Space Science* 139.1, s. 1–4.
-  Feltens, J. (2009). “Vector met hod to compute the Cartesian (X, Y, Z) to geodetic transformation on a triaxial ellipsoid”. *Journal of Geodesy* 83.2, s. 129–137.
-  Fukushima, T. (1999). “Fast transform from geocentric to geodetic coordinates”. *Journal of Geodesy* 73.11, s. 603–610.
-  Laskowski, P. (1991). “Is Newton’s iteration faster than simple iteration for transformation between geocentric and geodetic coordinates?” *Bulletin géodésique* 65.1, s. 14–17.

Wybrane algorytmy (kont.)



Ligas, M. i P. Banasik (2011). "Conversion between Cartesian and geodetic coordinates on a rotational ellipsoid by solving a system of nonlinear equations". *Geodesy and cartography* 60, s. 145–159.



Vermeille, H. (2002). "Direct transformation from geocentric coordinates to geodetic coordinates". *Journal of Geodesy* 76.8, s. 451–454.